

VU Research Portal

Prevention and reversal of atrial cardiomyocyte remodeling in Atrial Fibrillation

Hu, Xu

2019

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Hu, X. (2019). *Prevention and reversal of atrial cardiomyocyte remodeling in Atrial Fibrillation: via modulation of HSPs, metabolic and epigenetic factors*. [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam].

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Chapter 2

Keeping up the balance: Role of HDACs in cardiac proteostasis and therapeutic implications for atrial fibrillation

Deli Zhang¹, Xu Hu¹, Robert H. Henning¹, Bianca J.J.M. Brundel^{1,2#}

¹ Department of Clinical Pharmacy and Pharmacology, University Medical Centre Groningen, University of Groningen, The Netherlands.

² Department of Physiology, Institute for Cardiovascular Research, VU University Medical Center, The Netherlands

Abstract

Cardiomyocytes are long-lived post-mitotic cells with limited regenerative capacity. A proper cardiac function critically depends on the maintenance of a healthy proteostasis of protein expression, folding, assembly, trafficking, function and degradation, together commonly referred to as proteostasis. Impairment of proteostasis has a prominent role in the pathophysiology of aging-related neurodegenerative diseases, including Huntington's, Parkinson' and Alzheimer's disease. Emerging evidence reveals also a role for impaired proteostasis in the pathophysiology of common human cardiac diseases, such as cardiac hypertrophy, dilated and ischaemic cardiomyopathies, and AF. Histone deacetylases (HDACs) have been recently recognized as key modulators which control cardiac proteostasis by deacetylating various proteins. By deacetylating chromatin proteins, including histones, HDACs modulate epigenetic regulation of pathological gene expression. Also, HDACs exert a broad range of functions other than the nucleus by deacetylating structural and contractile proteins. The cytosolic actions of HDACs result in changed protein function through post-translational modifications and/or modulation of their degradation. Chapter 2 gives an overview of the mechanisms underlying the derailment of proteostasis in AF and subsequently focus on the role of HDACs herein. In addition, the therapeutic potential of HDAC inhibition to maintain a healthy proteostasis resulting a delay on AF onset and progression is discussed.

Keywords: Atrial Fibrillation, Histone deacetylases, Proteostasis, HDAC6

1.Introduction

1.1 Proteostasis

The proper function of cells depends critically on the maintenance of a healthy homeostasis of protein expression, folding, assembly, trafficking and degradation, together commonly referred to as proteostasis.^{1, 2} A proper proteostasis enables healthy cell and organismal development.¹ The requirement of maintenance of proteostasis is of importance, especially to long-lived post-mitotic cells, such as cardiomyocytes, that display limited regenerative capacity.^{3, 4} Derailment of proteostasis plays a prominent role in the pathophysiology of ageing-related neurodegenerative diseases, including Huntington's disease, Parkinson's disease, and Alzheimer's disease.⁴ The impairment of proteostasis is recently also implicated in the pathophysiology of common human cardiac diseases, including pathologic cardiac hypertrophy, dilated and ischemic cardiomyopathies and atrial fibrillation (AF).³⁻⁶

1.2 Remodeling in atrial fibrillation

AF is the most common progressive clinical tachyarrhythmia.⁷ Its incidence is age-related and increasing in the ageing population. AF can be caused by underlying cardiovascular conditions, including hypertension, cardiac surgery, pericarditis, congestive heart failure,^{8, 9} but clinical signs of underlying heart diseases are absent in about 30% of AF patients. However, additional risk factors are linked to this group of 'lone' AF patients, including alcohol abuse, obesity, metabolic syndrome, psychological stress and genetic factors.¹⁰⁻¹² AF-associated risk factors, but also AF itself, result in atrial arrhythmogenic remodeling, which is central to AF progression and defined as any change in atrial structure or function that promotes atrial arrhythmia.¹³ During the past decades, various mechanisms have been identified which promote the occurrence or maintenance of AF, including electrical remodeling and structural remodeling.¹⁴ Compared to electrical remodeling, structural remodeling is perceived as the main contributor to initiation and persistence of AF and may be present before the first episode of AF due to associated underlying diseases.⁵ Atrial structural remodeling, as observed in animal models and clinical AF, includes hypertrophy, dedifferentiation of cardiomyocytes and fibrosis. In addition, at the subcellular level, loss of contractile apparatus (myolysis) and changes in size and shape of the mitochondria, disruption of the sarcoplasmic reticulum and homogeneous distribution of nuclear heterochromatin are found.¹⁵⁻¹⁹ Although structural remodeling induces the vulnerability for AF, the molecular mechanisms underlying remodeling are not completely resolved.

1.3 Role of HDACs in AF associated derailment of proteostasis

Recent mechanistic studies show that the progressive derailment of proteostasis creates a substrate for the initiation of AF, with histone deacetylases (HDACs) playing a central role (Figure 1).^{3, 5, 16, 18, 20-24} HDACs have recently been recognized as key modulators in controlling cardiac proteostasis by changing acetylation status of various proteins.²⁵ By modifying chromatin they modulate epigenetic regulation of pathological gene expression. In addition, HDACs also exert a broad range of functions outside the nucleus by deacetylating structural and contractile proteins, leading to changes in protein function through post translational modifications (PTMs) and/or modulation of their degradation.^{6, 26-}

²⁸ Studies from the past decade indicate that HDACs alter proteostasis in cardiovascular diseases, including cardiac hypertrophy and heart failure, mainly by epigenetic regulation of pathological gene expression.²⁹⁻³⁵ In AF, however, involvement of HDACs in derailment of proteostasis seems much more associated with changes in PTM of contractile and structural proteins and the degradation of proteins.^{6, 28, 36} In the current review, we will summarize key mechanisms involved in the derailment of proteostasis and the modulating role of HDACs in AF progression. In addition, the potential of HDAC inhibition as a novel therapeutic target to treat clinical AF is discussed.




Figure 1: Overview of derailed proteostasis and the role of HDAC in AF progression. Risk factors for AF, such as hypertension, heart diseases, ischemia and diabetes, induce cardiac stress which results in derailment of proteostasis by impairment of protein expression, post translational modifications and degradation. Derailment of proteostasis is regulated by various HDACs, as indicated, and triggers structural remodeling in AF, such as cellular hypertrophy, dedifferentiation, myolysis and fibrosis. These structural changes create a substrate for further AF progression.

2. Central mechanisms of derailed proteostasis in AF

Recent studies revealed that three mechanisms importantly derail cardiomyocyte proteostasis in AF. These mechanisms include changes in protein expression, PTM and protein degradation.

2.1 Changes in protein expression by transcriptional reprogramming

Various studies show that AF induces changes in protein expression, which will challenge the proteostasis of cardiomyocytes (Figure 1). Changed levels in protein expression are caused at the gene transcriptional level by transcriptional reprogramming. A recent study of the atrial transcriptome in patients with AF identified a decreased expression of targets of several transcription factors related to inflammation, oxidative and cellular stress responses and changes in ion channel expression, indicating substantial transcriptional reprogramming in clinical AF.³⁷ Various additional studies in

patients with persistent AF and in tachypaced dogs, document changes in expression of proteins involved in calcium signaling, such as reduced L-type calcium channel alpha 1 (Cav1.2) and sarcoplasmic reticulum (SR) Ca²⁺-ATPase expression³⁸ and increased sodium calcium exchanger expression (NCX1) which was found to enhancing the Ca²⁺ leak from SR, resulting in promotion of clinical AF.³⁹

In addition, dedifferentiation of the atrial cardiomyocytes shifts the gene expression profile towards a more fetal phenotype as observed in both experimental and clinical AF.^{17, 40, 41} Dedifferentiation includes re-expression of β -myosin heavy chain (MHC), smooth muscle α -actin (α -SMA).⁴¹⁻⁴⁵ Together, these data show that the protein expression profile of cardiomyocytes is affected in AF, both by the regulation of specific genes and by a more general transcriptional reprogramming.

2.2 Alterations in post translational modifications

PTMs of proteins are critically important in the regulation of handling of proteins by the cardiomyocyte and thus determine the fitness of proteostasis. PTM refers to the covalent and generally enzymatic modification of proteins during or after protein biosynthesis.⁴⁶ Proteins are synthesized by ribosomes, where mRNA is translated into polypeptide chains, which may subsequently undergo PTM to form the mature protein product. PTMs, including (de)phosphorylation and (de)acetylation, alter the function of proteins and are important components of cell signaling.⁴⁷ Acetylation and deacetylation of proteins play an important role in pathological cardiac remodeling and are involved in cardiac diseases, including heart failure and AF.^{25, 28-36, 48-50} Protein phosphorylation has been studied in detail in AF and is recognized to underlie AF promotion in experimental and clinical AF.^{14, 51-54} Also protein acetylation is found to play an important role in AF progression. In particular, deacetylation of α -tubulin contributes to the development of a substrate for AF progression.⁶ Besides α -tubulin deacetylation, protein acetylation regulates fibrosis formation, electrical activation (via connexin 40) and calcium handling and thereby promotes atrial arrhythmogenesis and AF inducibility.^{24, 55} Thus, in addition to protein phosphorylation, the protein acetylation status now emerges as a key contributor to AF initiation and progression (Figure 1).

2.3 Derailment by protein degradation

By employing a goat model, Ausma et al. were the first to show that AF is associated with the degradation of the sarcomeric structure, named myolysis.⁵⁶ Later studies confirmed that myolysis is a general characteristic of atrial structural remodeling in patients with persistent AF.^{16, 18} There are indications that myolysis is caused by persistent activation of proteases, especially calpain 1.^{6, 16, 21, 57} In addition to protease activation, activation of auxiliary cellular protein degradation pathways, such as autophagy may drive AF progression. Autophagy is a pathway removing damaged or long-lived proteins and organelles. Controlled autophagy during (mild) cardiac stress conditions, such as nutrient deprivation, hypoxia and oxidative stress, supports cardiomyocyte survival. In contrast, stress-induced, excessive activation of autophagy causes derailment of cell proteostasis by degrading essential proteins and organelles, thereby triggering autophagic cell damage and death as found in mitral valve regurgitation and cardiac hypertrophy.^{58, 59} Interestingly, both cardiac conditions often trigger AF onset.⁶⁰ Ongoing research from our lab indicates that autophagy indeed contributes to the initiation and progression of AF.⁶¹

The findings indicate that derailment of proteostasis via changes in protein expression, PTM and degradation contributes importantly to AF progression. Interestingly, in the past years, HDACs emerge as a key player in cell homeostasis, as they control proteostasis by acting on both chromatin-related and non-chromatin-related pathways.

3.Role of HDACs in cardiac proteostasis

3.1 General role of HDACs

The fundamental processes of proteostasis are all subject to regulation by HDACs, including protein expression, proper PTM, trafficking and clearance. In total, 18 mammalian HDACs are encoded by distinct genes and grouped into four classes on the basis of similarity to yeast transcriptional repressors (Figure 2).⁶² Class I HDACs (HDACs 1, 2, 3 and 8) are related to yeast reduced potassium dependency-3 (Rpd3), class II HDACs (HDACs 4, 5, 6, 7, 9 and 10) to yeast histone deacetylase 1 (Hda1), and class III HDACs (sirtuin 1–7), usually named as sirtuins, to yeast Sir2. Class II HDACs are further divided into two subclasses, IIa (HDACs 4, 5, 7 and 9) and IIb (HDACs 6 and 10). Finally, HDAC11 lacks homology with yeast HDAC enzymes and belongs to a fourth class. Classes I, II and IV HDACs have a highly conserved zinc-dependent deacetylase domain and are referred as classical HDACs (Figure 2). The classical HDACs show also high similarity in structure, enzymatic function, subcellular localization and expression pattern. In contrast, class III HDACs (sirtuins) utilize nicotinamide adenine dinucleotide (NAD⁺) as a co-factor for catalytic activity and are because of this mechanism associated with aging and various cardiac diseases including metabolic and hypertrophic heart diseases.⁶³⁻⁶⁵ Since this review is focused on the classical HDACs (class I, II and IV, Figure 2), which are effective targets of the commonly used HDAC inhibitors, we excluded class III HDACs from this review.

In general, HDACs are enzymes that catalyze the removal of acetyl-groups from lysine residues from both nucleosomal histone tails and various non-histone proteins, thereby altering gene expression and cell function, respectively.⁶⁶ Exceptions are the nuclear IIa HDACs, especially HDAC4 and 5, which are abundant in the heart. Under normal conditions, class IIa HDACs bind to transcription factors, such as myocyte enhancer factor 2 (MEF2), and thereby repress their activity and downstream pathological fetal gene expression (Figure 3). In response to stress signals, class IIa HDACs are phosphorylated and undergo nuclear export, resulting in derepression of the downstream pathological fetal genes.³⁴ Class IIa HDACs regulate gene expression through recruitment of class I HDACs and by modulation of various transcription factors, including MEF2 and serum response factor (SRF) among others, resulting in cardiac hypertrophy and fibrosis (Table 1).^{24, 67-76}




Figure 2: Overview of the classical zinc dependent histone deacetylase (HDAC) superfamily. The classical HDAC family consists of members of the class I, IIa, IIb and IV HDACs. Black rectangles indicate the conserved HDAC catalytic domain. Numbers following the HDAC domain indicate the number of amino acids in human. MEF2-binding sites and binding sites for the 14-3-3 chaperone protein sites are marked by grey squares. S represents the serine residue which can be phosphorylated by kinases. HDAC6 has a zinc-finger ubiquitin-specific protease (ZnF-UBP) domain, which can bind to ubiquitin and plays an important role in autophagy. (Figure is modified from Haberland et al. Nat Rev Genet. 2009.)⁶⁶

Table 1: Overview of cardiac transcription factors (TGs) regulated by HDACs

TF	HDACs	Pathological pathways	References
MEF2	HDAC4	Heart failure, hypertrophy	63, 67, 68, 69
	HDAC5		
	HDAC7		
	HDAC9		
KLF4	HDAC2	Cardiac hypertrophy	70
YY1	HDAC2	hypertrophy	71, 72
	HDAC4		
	HDAC 5		
NKX 2.5	HDAC5	Heart development, NCX1 expression	73
Myocardin/ SRF	HDAC5	Differentiation of smooth muscle cells, hypertrophic arrhythmia	24, 74
NF-κB	HDAC1	Cardiac infarction	75
	HDAC5		




Figure 3: Effects of cardiac stress signals on HDACs in proteostasis regulation. Class IIa HDACs normally repress pathological cardiac gene expression. In response to stress signals, kinases, including protein kinase C (PKC), protein kinase D (PKD) and calcium/calmodulin-dependent kinase (CaMK), directly phosphorylate class IIa HDACs to trigger their nuclear export. The nuclear export of class IIa HDACs triggers derepression of downstream target pathological genes and thereby contributes to structural remodeling such as fibrosis, hypertrophy and changes of ion channel expression. In the cytoplasm, cardiac stress activates HDAC6, HDAC3 and HDAC4. HDAC6 deacetylates substrates such as α -tubulin and cortactin, resulting in degradation of structural proteins and functional loss. In addition, during stress, HDAC3²⁷, HDAC4²⁶ and HDAC6²⁸ colocalize with sarcomeric proteins and contribute to hypocontractility by deacetylation of sarcomeric and cytoskeletal proteins.

3.2 Role of HDACs in transcriptional reprogramming

The involvement of HDACs in AF is recently recognized. Two studies indicate that HDAC modulate transcriptional reprogramming in AF, possibly via epigenetic regulation. Liu et al.²⁴ were the first to show that HDAC inhibition in cardiac hypertrophy protects homeo-domain-only protein (HopX) transgenic mice from atrial arrhythmia inducibility and fibrosis.²⁴ In this study, HopX mice, which recruit HDAC to the chromatin to induce SRF-dependent transcription and cardiac hypertrophy, were either treated or untreated with the pan-HDAC inhibitor trichostatin A (TSA) and compared to control groups. TSA treated mice were protected against tachypacing-induced atrial arrhythmias and fibrosis formation. Moreover, this study suggests AF to originate from epigenetic regulation, since these mice feature increased HDAC activity, recruited by overexpressing the HopX. In addition, Lkhagva et al.

recently found that HDAC inhibition by the pan-HDAC inhibitor TSA and class I HDAC inhibitor MPTOE014, reduces arrhythmogenesis and AF inducibility by reducing calcium sparks via normalization of NCX1 and ryanodine receptor expression in rabbit pulmonary vein cardiomyocytes.⁵⁵ Also, HDACs may potentially modulate AF-induced activation of the fetal gene program via the chromatin-remodeling protein, Brg1. Brg1 preserves fetal cardiac differentiation by interacting with HDAC and poly (ADP ribose) polymerase (PARP), thereby repressing α -MHC and activating β -MHC expression.⁷⁷ Brg1 expression is turned off in adult cardiomyocytes and becomes reactivated during cardiac stress. Upon reactivation, Brg1 forms a complex with HDAC and PARP, in turn resulting in the α -MHC to β -MHC shift. Treatment of SW13 cells with TSA protects the cells from Brg1-induced shift of α -MHC to β -MHC.⁷⁷ Thus the apparent role of HDACs in AF warrants the exploration of the modulating role of HDACs in transcriptional reprogramming in AF-induced structural and functional cardiomyocyte remodeling.

3.3 HDAC modulate contractile function via PTM

Beyond altering chromatin structure and gene expression, HDAC also affect contractility by deacetylating contractile and structural proteins.^{6, 26-28} We recently identified a specific class IIb HDAC, HDAC6, to derail proteostasis via PTM of structural proteins, resulting in structural remodeling and AF promotion.⁶ Specifically, tachypacing increases HDAC6 expression and activity, resulting in deacetylation of α -tubulin in HL-1 cardiomyocytes and dog atrial cardiomyocytes. In turn, deacetylated α -tubulin is degraded by calpain, causing disruption of the microtubule structure and consequently contractile dysfunction.⁶ Importantly, comparable findings were obtained in patients with persistent AF, suggesting HDAC6-induced PTM via deacetylation and degradation of α -tubulin to represent a common feature of remodeling in both experimental and clinical AF. In addition to α -tubulin, also cortactin and HSP90 are reported as substrates of HDAC6.^{78, 79} Acetylated cortactin promotes polymerization of the actin cytoskeleton and regulates the structure of cardiomyocytes, actions that are counteracted by HDAC6 mediated deacetylation of cortactin. Under physiological conditions, HDAC6 complexes with HSP90 and heat shock factor-1 (HSF-1), thereby suppressing the heat shock response. Cardiac stress can induce the ubiquitination of proteins, which bind to the ubiquitin binding site of HDAC6, resulting in the dissociation of the complex. In turn, HSF-1 can translocate to the nucleus to induce the heat shock response, a response we previously found cardioprotective in AF.^{23, 80} Consequently, this action is independent of the deacetylation activity of HDAC6 and it is still unknown whether it plays a role in AF.

3.4 HDACs and protein degradation

As mentioned, AF induced HDAC6 activation initiates deacetylation of α -tubulin thus promoting a shift from the polymerized microtubule structure into its depolymerized form, thereby causing calcium transient loss and contractile dysfunction. Deacetylated and depolymerized α -tubulin is susceptible to accelerated degradation by the protease calpain.⁶ Also in patients with permanent AF, activation of HDAC6 deacetylates and degrades α -tubulin, and an inverse correlation with calpain activity is described. These results are consistent with previous reports demonstrating that microtubule network disruption causes changes in calcium signaling of cardiomyocytes, including reduction of contractions,⁸¹ L-type calcium current⁸² and calcium transient amplitude,⁸³ which are known to

underlie AF progression.⁵ Besides, the involvement of calpain is in accord with observations that calpain is strongly activated during experimental and clinical AF^{16, 18, 21} and with data showing degradation of brain α -tubulin by calpain.⁸⁴

3.5 Beyond AF – derailment of proteostasis and HDAC activation in other cardiac diseases

So far, HDAC6 was found to play a key role in experimental and clinical AF progression by derailment of cardiomyocyte proteostasis.⁶ In addition to its role in AF progression, HDAC6 is also involved in other cardiovascular diseases. McKinsey and colleagues observed that the catalytic activity of HDAC6 is consistently increased in myocardium subjected to hypertensive stimuli, but not during physiologic hypertrophy.⁵⁰ The HDAC6 catalytic activity is also induced by diverse extracellular stimuli, including norepinephrine and phenylephrine in cultured cardiomyocytes and fibroblasts.⁵⁰ Recently, they found that both HDAC6 null mice and mice treated with the HDAC6 inhibitor, tubastatin A, are protected against systolic dysfunction caused by angiotensin II. HDAC6 null mice also exhibited improved left ventricular function in the setting of pressure overload mediated by transverse aortic constriction.²⁸ HDAC6 inhibition appeared to preserve systolic function, in part, by enhancing myofibrillar force generation.²⁸ Furthermore, a very recent study showed that prevention of tubulin deacetylation by HDAC6 inhibition was protective in a mouse model of proteinopathy-induced heart failure.³⁶ Inhibiting α -tubulin deacetylation, by utilizing the FDA-approved drug suberoylanilide hydroxamic acid (SAHA), reduced protein aggregates in cardiomyocytes and resulted in a substantial improvement in cardiac function. Mechanistically, the inhibition of HDAC6 increases autophagy in cardiomyocytes, which can also be accomplished by voluntary exercise.³⁶ Taken together, next to the findings in AF, additional studies define novel roles for HDAC6 in various cardiac diseases and indicate the general therapeutic potential of HDAC6 inhibitors to treat cardiac dysfunction in patients.

4. Therapeutic implication for AF: HDAC6 as therapeutic target in AF

As we discussed above, derailment of proteostasis is involved in AF initiation and progression and HDAC6 represents an important modulator in cardiac proteostasis. Promoting maintenance of proteostasis by HDAC6 inhibition may prevent structural remodeling in cardiomyocytes and thereby attenuate AF progression.

Pan-HDAC inhibitors and class I HDAC inhibitors have been studied in various cardiac diseases and consistently reduced pathological cardiac remodeling, including heart failure.^{6, 6, 24, 25, 36, 50, 55, 64} In terms of potential toxicity and side effects, pan-HDAC inhibitors are generally regarded as effective and well tolerated for the treatment of cancer.⁸⁵ Reported side effects include nausea, fatigue, transient thrombocytopenia and, in some instances, myelosuppression.⁸⁶⁻⁸⁹ Since cancer therapy is frequently based on a maximum tolerated dose, the efficacious dose of pan-HDAC inhibitors in AF may be significantly lower and hence well tolerated. Among the HDACs, HDAC6 emerges as a key regulator in AF progression and therefore represents an interesting druggable target. Inhibition of HDAC6 by tubacin protects against AF-induced de-acetylation and subsequent depolymerization and degradation of α -tubulin by calpain.⁶ In this way, tubacin conserves α -tubulin proteostasis, and contractile function in experimental cardiomyocyte and *Drosophila* model systems for AF.⁶ However,

tubacin is not suitable for *in vivo* studies, because of its high lipophilicity (LogP = 6.36) and complicated synthesis.⁹⁰ Other HDAC6 inhibitors, such as tubastatin A and ricolinostat (ACY-1215) have been developed, and have shown beneficial effects in mouse models for neurodegenerative diseases and cancer.⁹⁰⁻⁹² Recently, the first evidence was provided for the efficacy of tubastatin A in the dog model for AF.⁶ Dogs treated with tubastatin A were protected against atrial tachypacing-induced derailment of α -tubulin homeostasis and electrical remodeling, resulting in improved cellular Ca^{2+} handling and contractile function, effectively counteracting AF progression. These *in vivo* findings strengthen the notion that HDAC6 inhibitors represent a novel therapeutic approach in AF. Trials with tubastatin A or ricolinostat should be initiated to determine the clinical potential of HDAC6 inhibitors in the treatment of first onset AF and the prevention of post-surgery AF. Interestingly, ricolinostat is currently tested in phase I and II clinical trials for the treatment of multiple myeloma, and so far, no serious side effects have been reported, making it an interesting candidate drug to test in patients with AF.⁹³

In summary, the proper function of cardiomyocytes depends critically on the maintenance of a healthy proteostasis. Impairment of proteostasis has a prominent role in the pathophysiology of common human cardiac diseases including AF. So far, HDAC6 has been recognized as key modulator controlling cardiac proteostasis by changing acetylation status of structural proteins, leading to changes in protein function and the modulation of their degradation. Therefore, inhibitors of HDAC6 may represent a novel therapeutic target to treat clinical AF.

Acknowledgements

This work was supported by the Dutch Heart Foundation (2009B024, 2013T096 and 2013T144) and the LSH-Impulse grant (40-43100-98-008).

References

1. Balch WE, Morimoto RI, Dillin A, Kelly JW. Adapting proteostasis for disease intervention. *Science* 2008;**319**:916-919.
2. Powers ET, Morimoto RI, Dillin A, Kelly JW, Balch WE. Biological and chemical approaches to diseases of proteostasis deficiency. *Annu Rev Biochem* 2009;**78**:959-991.
3. Meijering RA, Henning RH, Brundel BJ. Reviving the protein quality control system: therapeutic target for cardiac disease in the elderly. *Trends Cardiovasc Med* 2015;**25**:243-247.
4. Willis MS, Patterson C. Proteotoxicity and cardiac dysfunction--Alzheimer's disease of the heart? *N Engl J Med* 2013;**368**:455-464.
5. Meijering RA, Zhang D, Hoogstra-Berends F, Henning RH, Brundel BJ. Loss of proteostatic control as a substrate for atrial fibrillation: a novel target for upstream therapy by heat shock proteins. *Front Physiol* 2012;**3**:36.
6. Zhang D, Wu CT, Qi X, Meijering RA, Hoogstra-Berends F, Tadevosyan A, Cubukcuoglu Deniz G, Durdu S, Akar AR, Sibon OC, Nattel S, Henning RH, Brundel BJ. Activation of histone deacetylase-6 induces contractile dysfunction through derailment of alpha-tubulin proteostasis in experimental and human atrial fibrillation. *Circulation* 2014;**129**:346-358.
7. Dobrev D, Nattel S. New antiarrhythmic drugs for treatment of atrial fibrillation. *The Lancet* 2010;**375**:1212-1223.
8. January CT, Wann LS, Alpert JS, Calkins H, Cigarroa JE, Cleveland JC, Jr, Conti JB, Ellinor PT, Ezekowitz MD, Field ME, Murray KT, Sacco RL, Stevenson WG, Tchou PJ, Tracy CM, Yancy CW. 2014 AHA/ACC/HRS Guideline for the

Management of Patients With Atrial Fibrillation: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the Heart Rhythm Society. *Circulation* 2014;**130**:2071-2104.

9. Kourliouros A, Savelieva I, Kiotsekoglou A, Jahangiri M, Camm J. Current concepts in the pathogenesis of atrial fibrillation. *Am Heart J* 2009;**157**:243-252.
10. Fox CS, Parise H, D'Agostino RB S, Lloyd-Jones DM, Vasan RS, Wang TJ, Levy D, Wolf PA, Benjamin EJ. Parental atrial fibrillation as a risk factor for atrial fibrillation in offspring. *JAMA* 2004;**291**:2851-2855.
11. Saffitz JE. Connexins, conduction, and atrial fibrillation. *N Engl J Med* 2006;**354**:2712-2714.
12. Tsang TS, Miyasaka Y, Barnes ME, Gersh BJ. Epidemiological profile of atrial fibrillation: a contemporary perspective. *Prog Cardiovasc Dis* 2005;**48**:1-8.
13. Nattel S, Burstein B, Dobrev D. Atrial remodeling and atrial fibrillation: mechanisms and implications. *Circ Arrhythm Electrophysiol* 2008;**1**:62-73.
14. Nattel S, Harada M. Atrial remodeling and atrial fibrillation: recent advances and translational perspectives. *J Am Coll Cardiol* 2014;**63**:2335-2345.
15. De Jong AM, Maass AH, Oberdorf-Maass SU, Van Veldhuisen DJ, Van Gilst WH, Van Gelder IC. Mechanisms of atrial structural changes caused by stretch occurring before and during early atrial fibrillation. *Cardiovasc Res* 2011;**89**:754-765.
16. Brundel BJM, Ausma J, van Gelder IC, Van Der Want JJL, van Gilst WH, Crijns HJGM, Henning RH. Activation of proteolysis by calpains and structural changes in human paroxysmal and persistent atrial fibrillation. *Cardiovasc Res* 2002;**54**:380-389.
17. Ausma J, Borgers M. Dedifferentiation of atrial cardiomyocytes: from in vivo to in vitro. *Cardiovasc Res* 2002;**55**:9-12.
18. Ke L, Qi XY, Dijkhuis AJ, Chartier D, Nattel S, Henning RH, Kampinga HH, Brundel BJM. Calpain mediates cardiac troponin degradation and contractile dysfunction in atrial fibrillation. *J Mol Cell Cardiol* 2008;**45**:685-693.
19. Frustaci A, Chimenti C, Bellocci F, Morgante E, Russo MA, Maseri A. Histological substrate of atrial biopsies in patients with lone atrial fibrillation. *Circulation* 1997;**96**:1180-1184.
20. Ke L, Meijering RA, Hoogstra-Berends F, Mackovicova K, Vos MJ, Van Gelder IC, Henning RH, Kampinga HH, Brundel BJ. HSPB1, HSPB6, HSPB7 and HSPB8 protect against RhoA GTPase-induced remodeling in tachypaced atrial myocytes. *PLoS One* 2011;**6**:e20395.
21. Zhang D, Ke L, Mackovicova K, Van Der Want JJ, Sibon OC, Tanguay RM, Morrow G, Henning RH, Kampinga HH, Brundel BJ. Effects of different small HSPB members on contractile dysfunction and structural changes in a *Drosophila melanogaster* model for Atrial Fibrillation. *Journal of Molecular and Cellular Cardiology* 2011;**51**:381-9.
22. Brundel BJ, Ke L, Dijkhuis AJ, Qi X, Shiroshita-Takeshita A, Nattel S, Henning RH, Kampinga HH. Heat shock proteins as molecular targets for intervention in atrial fibrillation. *Cardiovasc Res* 2008;**78**:422-8.
23. Brundel BJ, Henning RH, Ke L, van Gelder IC, Crijns HJ, Kampinga HH. Heat shock protein upregulation protects against pacing-induced myolysis in HL-1 atrial myocytes and in human atrial fibrillation. *J Mol Cell Cardiol* 2006;**41**:555-62.
24. Liu F, Levin MD, Petrenko NB, Lu MM, Wang T, Yuan LJ, Stout AL, Epstein JA, Patel VV. Histone-deacetylase inhibition reverses atrial arrhythmia inducibility and fibrosis in cardiac hypertrophy independent of angiotensin. *J Mol Cell Cardiol* 2008;**45**:715-723.
25. McKinsey TA. Therapeutic potential for HDAC inhibitors in the heart. *Annu Rev Pharmacol Toxicol* 2012;**52**:303-319.
26. Gupta MP, Samant SA, Smith SH, Shroff SG. HDAC4 and PCAF bind to cardiac sarcomeres and play a role in regulating myofilament contractile activity. *J Biol Chem* 2008;**283**:10135-10146.

27. Samant SA, Courson DS, Sundaresan NR, Pillai VB, Tan M, Zhao Y, Shroff SG, Rock RS, Gupta MP. HDAC3-dependent reversible lysine acetylation of cardiac myosin heavy chain isoforms modulates their enzymatic and motor activity. *J Biol Chem* 2011;**286**:5567-5577.
28. Demos-Davies KM, Ferguson BS, Cavin MA, Mahaffey JH, Williams SM, Spiltoir JJ, Schuetz KB, Horn TR, Chen B, Ferrara C, Scellini B, Piroddi N, Tesi C, Poggesi C, Jeong MY, McKinsey TA. HDAC6 contributes to pathological responses of heart and skeletal muscle to chronic angiotensin-II signaling. *Am J Physiol Heart Circ Physiol* 2014;**307**:H252-8.
29. Ferguson BS, Harrison BC, Jeong MY, Reid BG, Wempe MF, Wagner FF, Holson EB, McKinsey TA. Signal-dependent repression of DUSP5 by class I HDACs controls nuclear ERK activity and cardiomyocyte hypertrophy. *Proc Natl Acad Sci U S A* 2013;**110**:9806-9811.
30. Bush EW, McKinsey TA. Targeting histone deacetylases for heart failure. *Expert Opin Ther Targets* 2009;**13**:767-784.
31. Olson EN, Backs J, McKinsey TA. Control of cardiac hypertrophy and heart failure by histone acetylation/deacetylation. *Novartis found Symp* 2006;**274**:3-12;
32. Vega RB, Harrison BC, Meadows E, Roberts CR, Papst PJ, Olson EN, McKinsey TA. Protein kinases C and D mediate agonist-dependent cardiac hypertrophy through nuclear export of histone deacetylase 5. *Mol Cell Biol* 2004;**24**:8374-8385.
33. McKinsey TA. The biology and therapeutic implications of HDACs in the heart. *Handb Exp Pharmacol* 2011;**206**:57-78.
34. Zhang CL, McKinsey TA, Chang S, Antos CL, Hill JA, Olson EN. Class II histone deacetylases act as signal-responsive repressors of cardiac hypertrophy. *Cell* 2002;**110**:479-488.
35. Antos CL, McKinsey TA, Dreitz M, Hollingsworth LM, Zhang CL, Schreiber K, Rindt H, Gorczynski RJ, Olson EN. Dose-dependent blockade to cardiomyocyte hypertrophy by histone deacetylase inhibitors. *J Biol Chem* 2003;**278**:28930-28937.
36. McLendon PM, Ferguson BS, Osinska H, Bhuiyan MS, James J, McKinsey TA, Robbins J. Tubulin hyperacetylation is adaptive in cardiac proteotoxicity by promoting autophagy. *Proc Natl Acad Sci U S A* 2014;**111**:E5178-86.
37. Deshmukh A, Barnard J, Sun H, Newton D, Castel L, Pettersson G, Johnston D, Roselli E, Gillinov AM, McCurry K, Moravec C, Smith JD, Van Wagoner DR, Chung MK. Left atrial transcriptional changes associated with atrial fibrillation susceptibility and persistence. *Circ Arrhythm Electrophysiol* 2015;**8**:32-41.
38. Brundel BJ, van Gelder IC, Henning RH, Tuinenburg AE, Deelman LE, Tieleman RG, Grandjean JG, van Gilst WH, Crijns HJ. Gene expression of proteins influencing the calcium homeostasis in patients with persistent and paroxysmal atrial fibrillation. *Cardiovasc Res* 1999;**42**:443-54.
39. Voigt N, Li N, Wang Q, Wang W, Trafford AW, Abu-Taha I, Sun Q, Wieland T, Ravens U, Nattel S, Wehrens XH, Dobrev D. Enhanced sarcoplasmic reticulum Ca²⁺ leak and increased Na⁺-Ca²⁺ exchanger function underlie delayed afterdepolarizations in patients with chronic atrial fibrillation. *Circulation* 2012;**125**:2059-2070.
40. Ausma J, Wijffels M, van Eys G, Koide M, Ramaekers F, Allessie M, Borgers M. Dedifferentiation of atrial cardiomyocytes as a result of chronic atrial fibrillation. *Am J Pathol* 1997;**151**:985-997.
41. Rucker-Martin C, Pecker F, Godreau D, Hatem SN. Dedifferentiation of atrial myocytes during atrial fibrillation: role of fibroblast proliferation in vitro. *Cardiovasc Res* 2002;**55**:38-52.
42. Cardin S, Libby E, Pelletier P, Le Bouter S, Shiroshita-Takeshita A, Le Meur N, Leger J, Demolombe S, Ponton A, Glass L, Nattel S. Contrasting gene expression profiles in two canine models of atrial fibrillation. *Circ Res* 2007;**100**:425-433.
43. Mace LC, Yermalitskaya LV, Yi Y, Yang Z, Morgan AM, Murray KT. Transcriptional remodeling of rapidly stimulated HL-1 atrial myocytes exhibits concordance with human atrial fibrillation. *J Mol Cell Cardiol* 2009;**47**:485-492.

44. Belus A, Piroddi N, Ferrantini C, Tesi C, Cazorla O, Toniolo L, Drost M, Mearini G, Carrier L, Rossi A, Mugelli A, Cerbai E, van der Velden J, Poggesi C. Effects of chronic atrial fibrillation on active and passive force generation in human atrial myofibrils. *Circ Res* 2010;**107**:144-152.
45. Mihm MJ, Yu F, Carnes CA, Reiser PJ, McCarthy PM, Van Wagoner DR, Bauer JA. Impaired myofibrillar energetics and oxidative injury during human atrial fibrillation. *Circulation* 2001;**104**:174-180.
46. Mann M, Jensen ON. Proteomic analysis of post-translational modifications. *Nat Biotechnol* 2003;**21**:255-261.
47. Yang X, Seto E. Lysine acetylation: codified crosstalk with other posttranslational modifications. *Mol Cell* 2008;**31**:449-461.
48. Williams SM, Golden-Mason L, Ferguson BS, Schuetze KB, Cavasin MA, Demos-Davies K, Yeager ME, Stenmark KR, McKinsey TA. Class I HDACs regulate angiotensin II-dependent cardiac fibrosis via fibroblasts and circulating fibrocytes. *J Mol Cell Cardiol* 2014;**67**:112-125.
49. Cavasin MA, Demos-Davies K, Horn TR, Walker LA, Lemon DD, Birdsey N, Weiser-Evans MC, Harral J, Irwin DC, Anwar A, Yeager ME, Li M, Watson PA, Nemenoff RA, Buttrick PM, Stenmark KR, McKinsey TA. Selective class I histone deacetylase inhibition suppresses hypoxia-induced cardiopulmonary remodeling through an antiproliferative mechanism. *Circ Res* 2012;**110**:739-748.
50. Lemon DD, Horn TR, Cavasin MA, Jeong MY, Haubold KW, Long CS, Irwin DC, McCune SA, Chung E, Leinwand LA, McKinsey TA. Cardiac HDAC6 catalytic activity is induced in response to chronic hypertension. *J Mol Cell Cardiol* 2011;**51**:41-50.
51. Neef S, Dybkova N, Sossalla S, Ort KR, Fluschnik N, Neumann K, Seipelt R, Schondube FA, Hasenfuss G, Maier LS. CaMKII-dependent diastolic SR Ca²⁺ leak and elevated diastolic Ca²⁺ levels in right atrial myocardium of patients with atrial fibrillation. *Circ Res* 2010;**106**:1134-1144.
52. Chelu MG, Sarma S, Sood S, Wang S, van Oort RJ, Skapura DG, Li N, Santonastasi M, Muller FU, Schmitz W, Schotten U, Anderson ME, Valderrabano M, Dobrev D, Wehrens XH. Calmodulin kinase II-mediated sarcoplasmic reticulum Ca²⁺ leak promotes atrial fibrillation in mice. *J Clin Invest* 2009;**119**:1940-1951.
53. Li D, Shinagawa K, Pang L, Leung TK, Cardin S, Wang Z, Nattel S. Effects of angiotensin-converting enzyme inhibition on the development of the atrial fibrillation substrate in dogs with ventricular tachypacing-induced congestive heart failure. *Circulation* 2001;**104**:2608-2614.
54. Goette A, Staack T, Rocken C, Arndt M, Geller JC, Huth C, Ansorge S, Klein HU, Lendeckel U. Increased expression of extracellular signal-regulated kinase and angiotensin-converting enzyme in human atria during atrial fibrillation. *J Am Coll Cardiol* 2000;**35**:1669-1677.
55. Lkhagva B, Chang S, Chen Y, Kao Y, Lin Y, Chiu CT, Chen S, Chen Y. Histone deacetylase inhibition reduces pulmonary vein arrhythmogenesis through calcium regulation. *Int J Cardiol* 2014;**177**:982-989.
56. Ausma J, Wijffels M, Thone F, Wouters L, Allessie M, Borgers M. Structural changes of atrial myocardium due to sustained atrial fibrillation in the goat. *Circulation* 1997;**96**:3157-3163.
57. Brundel BJ, Kampinga HH, Henning RH. Calpain inhibition prevents pacing-induced cellular remodeling in a HL-1 myocyte model for atrial fibrillation. *Cardiovasc Res* 2004;**62**:521-8.
58. Chen M, Chang J, Wang Y, Liu W, Ho W, Chang H. Autophagy as a mechanism for myolysis of cardiomyocytes in mitral regurgitation. *Eur J Clin Invest* 2011;**41**:299-307.
59. Gurusamy N, Das DK. Is autophagy a double-edged sword for the heart? *Acta Physiol Hung* 2009;**96**:267-276.
60. Zhu H, Tannous P, Johnstone JL, Kong Y, Shelton JM, Richardson JA, Le V, Levine B, Rothermel BA, Hill JA. Cardiac autophagy is a maladaptive response to hemodynamic stress. *J Clin Invest* 2007;**117**:1782-1793.
61. Meijering RA, Wiersma M, Deli Zhang F, Hoogstra-Berends XQ, Stanley Nattel RH, Henning BJ. Inhibition of ER stress-induced autophagy preserves proteostasis and protects against cardiomyocyte dysfunction in experimental and human Atrial Fibrillation. *Loss of Proteostasis as a Substrate for Atrial Fibrillation* 2014;:153-180 (thesis).

62. Gregoret IV, Lee YM, Goodson HV. Molecular evolution of the histone deacetylase family: functional implications of phylogenetic analysis. *J Mol Biol* 2004;**338**:17-31.
63. Ferguson BS, McKinsey TA. Non-sirtuin histone deacetylases in the control of cardiac aging. *J Mol Cell Cardiol* 2015;**83**:14-20.
64. Donmez G, Guarente L. Aging and disease: connections to sirtuins. *Aging Cell* 2010;**9**:285-290.
65. Pillai VB, Sundaresan NR, Gupta MP. Regulation of Akt signaling by sirtuins: its implication in cardiac hypertrophy and aging. *Circ Res* 2014;**114**:368-378.
66. Haberland M, Montgomery RL, Olson EN. The many roles of histone deacetylases in development and physiology: implications for disease and therapy. *Nat Rev Genet* 2009;**10**:32-42.
67. Nebbioso A, Manzo F, Miceli M, Conte M, Manente L, Baldi A, De Luca A, Rotili D, Valente S, Mai A, Uziel A, Gronemeyer H, Altucci L. Selective class II HDAC inhibitors impair myogenesis by modulating the stability and activity of HDAC-MEF2 complexes. *EMBO Rep* 2009;**10**:776-782.
68. Potthoff MJ, Wu H, Arnold MA, Shelton JM, Backs J, McAnally J, Richardson JA, Bassel-Duby R, Olson EN. Histone deacetylase degradation and MEF2 activation promote the formation of slow-twitch myofibers. *J Clin Invest* 2007;**117**:2459-2467.
69. Lu J, McKinsey TA, Nicol RL, Olson EN. Signal-dependent activation of the MEF2 transcription factor by dissociation from histone deacetylases. *Proc Natl Acad Sci U S A* 2000;**97**:4070-4075.
70. McKinsey TA, Zhang CL, Olson EN. Activation of the myocyte enhancer factor-2 transcription factor by calcium/calmodulin-dependent protein kinase-stimulated binding of 14-3-3 to histone deacetylase 5. *Proc Natl Acad Sci U S A* 2000;**97**:14400-14405.
71. Kee HJ, Kook H. Kruppel-like factor 4 mediates histone deacetylase inhibitor-induced prevention of cardiac hypertrophy. *J Mol Cell Cardiol* 2009;**47**:770-780.
72. Sucharov CC, Dockstader K, McKinsey TA. YY1 protects cardiac myocytes from pathologic hypertrophy by interacting with HDAC5. *Mol Biol Cell* 2008;**19**:4141-4153.
73. Glenn DJ, Wang F, Chen S, Nishimoto M, Gardner DG. Endothelin-stimulated human B-type natriuretic peptide gene expression is mediated by Yin Yang 1 in association with histone deacetylase 2. *Hypertension* 2009;**53**:549-555.
74. Chandrasekaran S, Peterson RE, Mani SK, Addy B, Buchholz AL, Xu L, Thiagarajan T, Kasiganesan H, Kern CB, Menick DR. Histone deacetylases facilitate sodium/calcium exchanger up-regulation in adult cardiomyocytes. *FASEB J* 2009;**23**:3851-3864.
75. Cao D, Wang Z, Zhang CL, Oh J, Xing W, Li S, Richardson JA, Wang DZ, Olson EN. Modulation of smooth muscle gene expression by association of histone acetyltransferases and deacetylases with myocardin. *Mol Cell Biol* 2005;**25**:364-376.
76. Zhang LX, Zhao Y, Cheng G, Guo TL, Chin YE, Liu PY, Zhao TC. Targeted deletion of NF-kappaB p50 diminishes the cardioprotection of histone deacetylase inhibition. *Am J Physiol Heart Circ Physiol* 2010;**298**:H2154-63.
77. Hang CT, Yang J, Han P, Cheng HL, Shang C, Ashley E, Zhou B, Chang CP. Chromatin regulation by Brg1 underlies heart muscle development and disease. *Nature* 2010;**466**:62-67.
78. Zhang X, Yuan Z, Zhang Y, Yong S, Salas-Burgos A, Koomen J, Olashaw N, Parsons JT, Yang XJ, Dent SR, Yao TP, Lane WS, Seto E. HDAC6 modulates cell motility by altering the acetylation level of cortactin. *Mol Cell* 2007;**27**:197-213.
79. Boyault C, Zhang Y, Fritah S, Caron C, Gilquin B, Kwon SH, Garrido C, Yao TP, Vourc'h C, Matthias P, Khochbin S. HDAC6 controls major cell response pathways to cytotoxic accumulation of protein aggregates. *Genes Dev* 2007;**21**:2172-81.
80. Brundel BJ, Shiroshita-Takeshita A, Qi X, Yeh YH, Chartier D, van Gelder IC, Henning RH, Kampinga HH, Nattel S. Induction of heat shock response protects the heart against atrial fibrillation. *Circ Res* 2006;**99**:1394-402.
81. Kolodney MS, Elson EL. Contraction due to microtubule disruption is associated with increased phosphorylation of myosin regulatory light chain. *Proc Natl Acad Sci U S A* 1995;**92**:10252-10256.

82. Galli A, DeFelice LJ. Inactivation of L-type Ca channels in embryonic chick ventricle cells: dependence on the cytoskeletal agents colchicine and taxol. *Biophys J* 1994;**67**:2296-2304.
83. Leach R, Desai J, Orchard C. Effect of cytoskeleton disruptors on L-type Ca channel distribution in rat ventricular myocytes. *Cell Calcium* 2005;**38**:515-526.
84. Billger M, Wallin M, Karlsson JO. Proteolysis of tubulin and microtubule-associated proteins 1 and 2 by calpain I and II. Difference in sensitivity of assembled and disassembled microtubules. *Cell Calcium* 1988;**9**:33-44.
85. Marks PA. The clinical development of histone deacetylase inhibitors as targeted anticancer drugs. *Expert Opin Investig Drugs* 2010;**19**:1049-1066.
86. Galli M, Salmoiraghi S, Golay J, Gozzini A, Crippa C, Pescosta N, Rambaldi A. A phase II multiple dose clinical trial of histone deacetylase inhibitor ITF2357 in patients with relapsed or progressive multiple myeloma. *Ann Hematol* 2010;**89**:185-190.
87. Giles F, Fischer T, Cortes J, Garcia-Manero G, Beck J, Ravandi F, Masson E, Rae P, Laird G, Sharma S, Kantarjian H, Dugan M, Albitar M, Bhalla K. A phase I study of intravenous LBH589, a novel cinnamic hydroxamic acid analogue histone deacetylase inhibitor, in patients with refractory hematologic malignancies. *Clin Cancer Res* 2006;**12**:4628-4635.
88. Klimek VM, Fircanis S, Maslak P, Guernah I, Baum M, Wu N, Panageas K, Wright JJ, Pandolfi PP, Nimer SD. Tolerability, pharmacodynamics, and pharmacokinetics studies of depsipeptide (romidepsin) in patients with acute myelogenous leukemia or advanced myelodysplastic syndromes. *Clin Cancer Res* 2008;**14**:826-832.
89. O'Connor OA, Heaney ML, Schwartz L, Richardson S, Willim R, MacGregor-Cortelli B, Curly T, Moskowitz C, Portlock C, Horwitz S, Zelenetz AD, Frankel S, Richon V, Marks P, Kelly WK. Clinical experience with intravenous and oral formulations of the novel histone deacetylase inhibitor suberoylanilide hydroxamic acid in patients with advanced hematologic malignancies. *J Clin Oncol* 2006;**24**:166-173.
90. Butler KV, Kalin J, Brochier C, Vistoli G, Langley B, Kozikowski AP. Rational design and simple chemistry yield a superior, neuroprotective HDAC6 inhibitor, tubastatin A. *J Am Chem Soc* 2010;**132**:10842-6.
91. Santo L, Hideshima T, Kung AL, Tseng JC, Tamang D, Yang M, Jarpe M, van Duzer JH, Mazitschek R, Ogier WC, Cirstea D, Rodig S, Eda H, Scullen T, Canavese M, Bradner J, Anderson KC, Jones SS, Raje N. Preclinical activity, pharmacodynamic, and pharmacokinetic properties of a selective HDAC6 inhibitor, ACY-1215, in combination with bortezomib in multiple myeloma. *Blood* 2012;**119**:2579-2589.
92. d'Ydewalle C, Krishnan J, Chiheb DM, Van Damme P, Irobi J, Kozikowski AP, Berghe PV, Timmerman V, Robberecht W, Van Den Bosch L. HDAC6 inhibitors reverse axonal loss in a mouse model of mutant HSPB1-induced Charcot-Marie-Tooth disease. *Nat Med* 2011;**17**:968-974.
93. Ashjian E, Redic K. Multiple myeloma: Updates for pharmacists in the treatment of relapsed and refractory disease. *J Oncol Pharm Pract* 2015;.